



JPW

Dkt. 68262-A/JPW/PJP

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
For : **MESENCHYMAL STEM CELLS AS A VEHICLE FOR ION
CHANNEL TRANSFER**

1185 Avenue of the Americas
New York, New York 10036
May 14, 2004

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

SIR:

INFORMATION DISCLOSURE STATEMENT

In compliance with their duty of disclosure under 37 C.F.R. §1.56, applicants direct the Examiner's attention to the following references, which are listed on the accompanying form PTO-1449 as Exhibit 1.

The information identified on the attached form PTO-1449 was cited by or submitted to the Patent Office in prior application Serial No. 10/342,506 filed on January 15, 2003 which is relied upon for a filing date under 35 U.S.C. §120. Therefore, pursuant to 37 C.F.R. §1.98(d) copies of the information identified herein are not required.

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 2

1. DiFrancesco D: The cardiac hyperpolarizing-activated current, I_f : Origins and developments. *Prog. BiophysMol. Biol.* Vol. 46, No. 3, 1985, pages 163-183;
2. Zhou Z and Lipsius SL: Effect of isoprenaline on I_f current in latent pacemaker cells isolated from cat right atrium: ruptured vs. perforated patch whole-cell recording methods. *Pflugers Arch.* Vol. 423, No.5 Pt. 6, June 1993, pages 442-447;
3. Thuringer D, et al.: A hyperpolarization-activated inward current in human myocardial cells. *JmolCell. Cardiol.* Vol. 24, No. 5, May 1992, pages 451-455;
4. Rosen, M.R. and Robinson R.B.: Heart rate: a simple yet complex concept. *Dialogues in Cardiovascular Medicine.* Vol. 6, No. 1, 2001, pages 2-19;
5. Yu H, et al.: Pacemaker current exists in ventricular myocytes. *Circ. Res.* Vol. 72, No. 1, January 1993, pages 232-236.;
6. Cerbai E, et al.: The properties of the pacemaker current I_f in Human Ventricular Myocytes are modulaed by Cardiac Disease: *Jmol. Cell Cardiol.* Vol. 33, No. 3, March 2001, pages 441-448;
7. Yu, H. et al.: MinK-Related Peptide 1: A β Subunit for the HCN Ion Channel Subunit Family Enhances Expression and Speeds Activation. *Circ. Res.* Vol. 88, 2001, pages 84-87;
8. Robinson RB, et al.: Developmental change in the voltage dependence of the pacemaker current, I_f , in rat ventricle

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 3

- cells. *Pflugers Arch.* Vol. 433, 1991, pages 533-535;
9. Fares N, et al.: Characterization of a hyperpolarization-activated current in dedifferentiated adult rat ventricular cells in primary culture. *J. Physiol.* Vol. 506, No. 1, January 1, 1998, pages 73-82;
 10. Cerbai E, et al.: Influence of postnatal-development on I_f occurrence and properties in neonatal rat ventricular myocytes. *Cardiovasc. Res.* Vol. 42, No. 2, May 1999, pages 416-423;
 11. Cerbai E, et al.: Characterization of the hyperpolarization-activated current, I_f , in ventricular myocytes isolated from hypertensive rats. *J. Physiol.* Vol. 481, No. 3, Dec 15, 1994, pages 585-591;
 12. Cerbai E, et al.: Characterization of the hyperpolarization-activated current, I_f , in ventricular myocytes from human failing heart. *Circulation.* Vol. 95, No. 3, February 4, 1997, pages 568-571;
 13. Santoro B, et al.: Interactive cloning with the SH3 domain of N-src identifies a new brain specific ion channel protein, with homology to Eag and cyclic nucleotide-gated channels. *Proc. Natl. Sci. USA.* Vol. 94, No. 26, December 23, 1997, pages 14815-14820;
 14. Ludwig A, et al.: A family of hyperpolarization-activated mammalian cation channels. *Nature.* Vol. 393, No. 6685, June 11, 1998, pages 587-591;
 15. Santoro B, et al.: Identification of a gene encoding a

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 4

- hyperpolarization-activated pacemaker channel of brain.
Cell. Vol. 93, No. 5, May 29, 1998, pages 717-729;
16. Shi W, et al.: Distribution and Prevalence of hyperpolarization-activated cation channel (HCN) mRNA Expression in Cardiac Tissues. *Circ. Res.* Vol. 85, No. 1, July 9, 1999, pages e1-e6;
 17. Ishii TM, et al.: Molecular characterization of the hyperpolarization-activated cation channel in rabbit heart sinoatrial node. *J. Biol. Chem.* Vol. 264, No. 18, April 30, 1999, pages 12835-12839;
 18. Ludwig A, et al.: Two pacemaker channels from human heart with profoundly different activation kinetics. *EMBO J.* Vol. 18, No. 9, May 4, 1999, pages 2323-2329;
 19. Moosmang S, et al.: Cellular expression and functional characterization of four hyperpolarization-activated pacemaker channels in cardiac and neuronal tissues. *Eur. J. Biochem.* Vol. 268, No. 6, March 2001, pages 1646-1652;
 20. Qu, J. et al.: Functional Comparison of HCN isoforms expressed in ventricular and HEK 293 cells. *Pflügers Arch. - Eur. J. Physiol.* Vol. 444, 2002, pages 597-601;
 21. Protas L, et al.: Chronic neuropeptide Y exposure increases L-type Ca current in neonatal rat cardiomyocytes. *Am. J. Physiol.* Vol. 277, No. 3 Pt. 2, September 1999, pages H940-H946;
 22. Kuznetsov V, et al.: β 2-adrenergic receptor actions in neonatal and adult rat ventricular myocytes. *Circ. Res.* Vol.

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 5

- 76, No. 1, January 1995, pages 40-52;
23. Qu, J. et al.: Sympathetic innervation alters activation of pacemaker current (I_f) in rat ventricle. *J. of Physiol.* Vol.526, No.3, 2000, pages 561-569;
24. Ng P, et al.: An enhanced system for construction of adenoviral vectors by the two-plasmid rescue method. *Hwn.Gene Ther.* Vol. 11, No. 5, March 20, 2000, pages 693-699;
25. He TC, et al.: A simplified system for generating recombinant adenoviruses. *Proc. Natl. Acad. Sci. USA.* Vol. 95, No. 5, March 3, 1998, pages 2509-2514;
26. Santoro B, et al.: The HCN gene family: molecular basis of the hyperpolarization-activated pacemaker channels. *Ann. NY Acad. Sci.* Vol.868, April 30, 1999, pages 741-764;
27. Hamm, A. et al.: Efficient Transfection Method for Primary Cells. *Tissue Engineering.* Vol. 8, No.2, 2002, pages 235-245;
28. Hansen JE, et al.: Prediction of O-glycosylation of mammalian proteins:Specificity patterns of UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase. *Biochem. J.* Vol. 308, No. 3, June 15, 1995, pages 801-813;
29. Cui J, et al.: Gating of IsK expressed in *Xenopus* oocytes depends on the amount of mRNA injected. *Gen. Physiol.* Vol. 104, No. 1, July 1994, pages 87-105;
30. Guillemare E, et al.: Effects of the level of mRNA expression on biophysical properties, sensitivity to

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 6

- neurotoxins, and regulation of the brain delayed-rectifier K⁺ channels Kv1.2. *Biochemistry*. Vol. 31, No. 49, December 15, 1992, pages 12463-12468;
31. Honore E, et al.: Different types of K⁺ channel current are generated by different levels of a single mRNA. *EMBO J*. Vol. 11, No. 7, July 1992, pages 2465-2471;
 32. Qu, J. et al.: HCN2 Overexpression in Newborn and Adult Ventricular Myocytes. *Circ. Res*. Vol.89, 2001, pages 8-14;
 33. DiFrancesco D, et al.: Direct activation of cardiac pacemaker channels by intracellular cyclic AMP. *Nature*. Vol. 351, No. 6322, May 9, 1991, pages 145-147;
 34. Kaupp UB, et al.: Molecular diversity of pacemaker ion channels. *Annu, Rev. Physiol*. Vol. 63, 2001, pages 235-257;
 35. Chang F, et al.: Effects of protein kinase inhibitors on canine Purkinje fibre pacemaker depolarization and the pacemaker current I_f. *J. Physiol*. Vol. 440, 1991, pages 367-384;
 36. Yu H, et al.: Phosphatase inhibition by calyculin A increases I_f in canine Purkinje fibers and myocytes. *Pflugers Arch*. Vol. 422, No. 6, March 1993, pages 614-616;
 37. Schagger H and von Jagow G: Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for separation of proteins in the range from 1 to 100 Kda. *Analytical Biochem*. Vol. 166, No. 2, November 1, 1987, pages 368-379;
 38. Ranjan R, et al.: Mechanism of anode break stimulation in

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 7

- the heart. *Biophys. J.* Vol. 74, No. 4, April 1998, pages 1850-1863;
39. Moroni A, et al.: Kinetic and ionic properties of the human HCN2 pacemaker channel. *Pflugers Arch.* Vol. 439, No. 5, March 2000, pages 618-626;
40. Santoro B, et al.: Molecular and functional heterogeneity of hyperpolarization-activated pacemaker channels in the mouse CNS. *Jneurosci.* Vol. 20, No. 14, July 15, 2000, pages 5264-5275;
41. Valiunas, V.: Biophysical properties of connexin-45 gap junction hemichannels studied in vertebrate cells. *J. Gen. Physiol.* Vol.119, No.2, 2002, pages 147-164;
42. Melman YF, et al.: Structural determinants of KvLQT1 control by the KCNE family of proteins. *J Biol Chem.* Vol. 276, No. 9, March 2, 2001, pages 6439-6444;
43. Tinel N, et al.: KCNE2 confers background current characteristics to the cardiac KCNQ1 potassium channel. *EMBO J.* Vol. 19, No. 23, December 1, 2000, pages 6326-6330;
44. Martens JR, et al.: Differential targeting of Shaker-like potassium channels to lipid rafts. *BiolChem.* Vol. 275, No. 11, March 17, 2000, pages 7443-7446;
45. Chauhan VS, et al.: Abnormal cardiac Na(+) channel properties and QT heart rate adaptation in neonatal ankyrin(B) knockout mice. *Circ. Res.* Vol. 86, No. 4, March 3, 2000, pages 441-447;
46. Cao, F. et al.: A quantitative analysis of connexin-specific

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 8

- permeability differences of gap junctions expressed in HeLa transfectants and *Xenopus* oocytes. *J. Cell Sci.* Vol. 111 (pt.1), 1998, pages 31-43;
47. Gerhardstein BL, et al.: Proteolytic processing of the C terminus of the alpha (1C) subunit of L-type calcium channels and role of a proline-rich domain in membrane tethering of proteolytic fragments. *J Biol. Chem.* Vol. 275, No. 12, March 24, 2000, pages 8556-8563;
48. Barbuti A, et al.: Action of internal pronase on the f-channel kinetics in the rabbit SA node. *J. Physiol.* Vol. 520, No. 3, November 1, 1999, pages 737-744;
49. Wahler GM: Developmental increases in the inwardly rectifying potassium current of rat ventricular myocytes. *Am. J. Physiol.* Vol. 262, No. 5 Pt. 1, May 1992, pages C1266;
50. Sanguinetti MC, et al.: Coassembly of KvLGQT1 and minK (I_{SK}) proteins to form cardiac I_{SK} potassium channels. *Nature.* Vol. 384, No. 6604, November 7, 1996, pages 80-83;
51. Dixon JE and McKinnon D: Quantitative analysis of potassium channel expression in atrial and ventricular muscle of rats. *Circ. Res.* Vol. 75, No. 2, August 1994, pages 252-260;
52. Abbott GW, et al.: MiRP1 forms I_{kr} potassium channels with HERG and is associated with cardiac arrhythmia. *Cell.* Vol. 97, No. 2, April 16, 1999, pages 175-187;
53. Altomare C, et al.: Integrated allosteric model of voltage gating of HCN channels. *J. Gen. Physiol.* Vol. 117, No.6,

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 9

2001; pages 519-532;

54. Valiunas, V. et al.: Cardiac gap junction channels show quantitative differences in selectivity. *Circ. Res.* Vol.91, No.2, 2002, pages 104-111;
55. Accili EA, et al.: Properties and modulation of I_f in newborn versus adult cardiac SA node. *Am. J. Physiol.* Vol. 272, 1991, pages H1549-H1552;
56. Accili EA, et al.: Differential control of the hyperpolarization-activated current (I_f) by intracellular cAMP and phosphatase inhibition. *J. Physiol.* Vol. 491, 1996, pages 643-651;
57. Wainger BJ, et al.: Molecular Mechanism of cAMP modulation of HCN pacemaker channels. *Nature.* Vol.411, No. 6839, 2001 pages 805-810; and
58. Ellingsen O, et al.: Adult rat ventricular myocytes cultured in defined medium: phenotype and electromechanical function. *Am. J. Physiol.* Vol. 265, No. 2 Pt. 2, August 1993, pages H747-H754.

If a telephone interview would be of assistance in advancing prosecution of the subject application, applicants' undersigned attorney invites the Examiner to telephone him at the number provided below.

Applicants : Peter Brink et al.
Serial No. : 10/757,827
Filed : January 15, 2004
Page 10

Applicants are filing this Information Disclosure Statement under 37 C.F.R. '1.97(b)(3) before the mailing of a first Office Action on the merits. Accordingly, no fee is deemed necessary in connection with the filing of this Information Disclosure Statement. However, if a fee is required, authorization is hereby given to charge the amount of any such fee to Deposit Account No. 03-3125.

Respectfully submitted,



John P. White
Registration No. 28,678
Peter J. Phillips
Registration No. 29,691
Attorney for Applicant
Cooper & Dunham LLP
1185 Avenue of the Americas
New York, New York 10036
(212) 278-0400


I hereby certify that this correspondence is being deposited this date with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to:

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450



5/14/04

Peter J. Phillips Date
Reg. No. 29,691



Form PTO-1449 U.S. Department of Commerce Patent and Trademark Office INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)	Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
	Applicant Peter Brink et al.	
	Filing Date January 15, 2004	Group

U.S. PATENT DOCUMENTS

Examiner Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate

FOREIGN PATENT DOCUMENTS

	Document Number	Date	Country	Class	Subclass	Translation	
						Yes	No

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)

1	DiFrancesco D: The cardiac hyperpolarizing-activated current, I_f : Origins and developments. <i>Prog. BiophysMol. Biol.</i> Vol. 46, No. 3, 1985, pages 163-183;
2	Zhou Z and Lipsius SL: Effect of isoprenaline on I_f current in latent pacemaker cells isolated from cat right atrium: ruptured vs. perforated patch whole-cell recording methods. <i>Pflugers Arch.</i> Vol. 423, No.5 Pt. 6, June 1993, pages 442-447;
3	Thuringer D, et al.: A hyperpolarization-activated inward current in human myocardial cells. <i>JmolCell. Cardiol.</i> Vol. 24, No. 5, May 1992, pages 451-455;

EXAMINER	DATE CONSIDERED
----------	-----------------

*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.

Appl. : Peter Brink et al.
Serial #: 10/757,827
Filed : January 15, 2004
Exhibit 1

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	4	Rosen, M.R. and Robinson R.B.: Heart rate: a simple yet complex concept. <i>Dialogues in Cardiovascular Medicine</i> . Vol. 6, No. 1, 2001, pages 2-19;			
	5	Yu H, et al.: Pacemaker current exists in ventricular myocytes. <i>Circ. Res.</i> Vol. 72, No. 1, January 1993, pages 232-236.;			
	6	Cerbai E, et al.: The properties of the pacemaker current I_f in Human Ventricular Myocytes are modulated by Cardiac Disease. <i>Jmol. Cell Cardiol.</i> Vol. 33, No. 3, March 2001, pages 441-448;			
	7	Yu, H. et al.: MinK-Related Peptide 1: A β Subunit for the HCN Ion Channel Subunit Family Enhances Expression and Speeds Activation. <i>Circ. Res.</i> Vol. 88, 2001, pages 84-87;			
	8	Robinson RB, et al.: Developmental change in the voltage dependence of the pacemaker current, I_f , in rat ventricle cells. <i>Pflugers Arch.</i> Vol. 433, 1991, pages 533-535;			
	9	Fares N, et al.: Characterization of a hyperpolarization-activated current in dedifferentiated adult rat ventricular cells in primary culture. <i>J. Physiol.</i> Vol. 506, No. 1, January 1, 1998, pages 73-82;			
	10	Cerbai E, et al.: Influence of postnatal-development on I_f occurrence and properties in neonatal rat ventricular myocytes. <i>Cardiovasc. Res.</i> Vol. 42, No. 2, May 1999, pages 416-423;			
	11	Cerbai E, et al.: Characterization of the hyperpolarization-activated current, I_f , in ventricular myocytes isolated from hypertensive rats. <i>J. Physiol.</i> Vol. 481, No. 3, Dec 15, 1994, pages 585-591;			
	12	Cerbai E, et al.: Characterization of the hyperpolarization-activated current, I_f , in ventricular myocytes from human failing heart. <i>Circulation.</i> Vol. 95, No. 3, February 4, 1997, pages 568-571;			
	13	Santoro B, et al.: Interactive cloning with the SH3 domain of N-src identifies a new brain specific ion channel protein, with homology to Eag and cyclic nucleotide-gated channels. <i>Proc. Natl. Sci. USA.</i> Vol. 94, No. 26, December 23, 1997, pages 14815-14820;			
	14	Ludwig A, et al.: A family of hyperpolarization-activated mammalian cation channels. <i>Nature.</i> Vol. 393, No. 6685, June 11, 1998, pages 587-591;			
EXAMINER		DATE CONSIDERED			
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	15	Santoro B, et al.: Identification of a gene encoding a hyperpolarization-activated pacemaker channel of brain. <i>Cell</i> . Vol. 93, No. 5, May 29, 1998, pages 717-729;			
	16	Shi W, et al.: Distribution and Prevalence of hyperpolarization-activated cation channel (HCN) mRNA Expression in Cardiac Tissues. <i>Circ. Res.</i> Vol. 85, No. 1, July 9, 1999, pages e1-e6;			
	17	Ishii TM, et al.: Molecular characterization of the hyperpolarization-activated cation channel in rabbit heart sinoatrial node. <i>J. Biol. Chem.</i> Vol. 264, No. 18, April 30, 1999, pages 12835-12839;			
	18	Ludwig A, et al.: Two pacemaker channels from human heart with profoundly different activation kinetics. <i>EMBO J.</i> Vol. 18, No. 9, May 4, 1999, pages 2323-2329;			
	19	Moosmang S, et al.: Cellular expression and functional characterization of four hyperpolarization-activated pacemaker channels in cardiac and neuronal tissues. <i>Eur. J. Biochem.</i> Vol. 268, No. 6, March 2001, pages 1646-1652;			
	20	Qu, J. et al.: Functional Comparison of HCN isoforms expressed in ventricular and HEK 293 cells. <i>Pfuglers Arch. - Eur. J. Physiol.</i> Vol. 444, 2002, pages 597-601;			
	21	Protas L, et al.: Chronic neuropeptide Y exposure increases L-type Ca current in neonatal rat cardiomyocytes. <i>Am. J. Physiol.</i> Vol. 277, No. 3 Pt. 2, September 1999, pages H940-H946;			
	22	Kuznetsov V, et al.: β 2-adrenergic receptor actions in neonatal and adult rat ventricular myocytes. <i>Circ. Res.</i> Vol. 76, No. 1, January 1995, pages 40-52;			
	23	Qu, J. et al.: Sympathetic innervation alters activation of pacemaker current (I_f) in rat ventricle. <i>J. of Physiol.</i> Vol. 526, No. 3, 2000, pages 561-569;			
	24	Ng P, et al.: An enhanced system for construction of adenoviral vectors by the two-plasmid rescue method. <i>Hum. Gene Ther.</i> Vol. 11, No. 5, March 20, 2000, pages 693-699;			
	25	He TC, et al.: A simplified system for generating recombinant adenoviruses. <i>Proc. Natl. Acad. Sci. USA.</i> Vol. 95, No. 5, March 3, 1998, pages 2509-2514;			
EXAMINER			DATE CONSIDERED		
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	26	Santoro B, et al.: The HCN gene family: molecular basis of the hyperpolarization-activated pacemaker channels. <i>Ann. NY Acad. Sci.</i> Vol.868, April 30, 1999, pages 741-764;			
	27	Hamm, A. et al.: Efficient Transfection Method for Primary Cells. <i>Tissue Engineering</i> . Vol. 8, No.2, 2002, pages 235-245;			
	28	Hansen JE, et al.: Prediction of O-glycosylation of mammalian proteins: Specificity patterns of UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase. <i>Biochem. J.</i> Vol. 308, No. 3, June 15, 1995, pages 801-813;			
	29	Cui J, et al.: Gating of Isk expressed in Xenopus oocytes depends on the amount of mRNA injected. <i>Gen. Physiol.</i> Vol. 104, No. 1, July 1994, pages 87-105;			
	30	Guillemare E, et al.: Effects of the level of mRNA expression on biophysical properties, sensitivity to neurotoxins, and regulation of the brain delayed-rectifier K ⁺ channels Kv1.2. <i>Biochemistry</i> . Vol. 31, No. 49, December 15, 1992, pages 12463-12468;			
	31	Honore E, et al.: Different types of K ⁺ channel current are generated by different levels of a single mRNA. <i>EMBO J.</i> Vol. 11, No. 7, July 1992, pages 2465-2471;			
	32	Qu, J. et al.: HCN2 Overexpression in Newborn and Adult Ventricular Myocytes. <i>Circ. Res.</i> Vol.89, 2001, pages 8-14;			
	33	DiFrancesco D, et al.: Direct activation of cardiac pacemaker channels by intracellular cyclic AMP. <i>Nature</i> . Vol. 351, No. 6322, May 9, 1991, pages 145-147;			
	34	Kaupp UB, et al.: Molecular diversity of pacemaker ion channels. <i>Annu. Rev. Physiol.</i> Vol. 63, 2001, pages 235-257;			
	35	Chang F, et al.: Effects of protein kinase inhibitors on canine Purkinje fibre pacemaker depolarization and the pacemaker current I _f . <i>J. Physiol.</i> Vol. 440, 1991, pages 367-384;			
	36	Yu H, et al.: Phosphatase inhibition by calyculin A increases I _f in canine Purkinje fibers and myocytes. <i>Pflugers Arch.</i> Vol. 422, No. 6, March 1993, pages 614-616;			
EXAMINER			DATE CONSIDERED		
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	37	Schagger H and von Jagow G: Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for separation of proteins in the range from 1 to 100 Kda. <i>Analytical Biochem.</i> Vol. 166, No. 2, November 1, 1987, pages 368-379;			
	38	Ranjan R, et al.: Mechanism of anode break stimulation in the heart. <i>Biophys. J.</i> Vol. 74, No. 4, April 1998, pages 1850-1863;			
	39	Moroni A, et al.: Kinetic and ionic properties of the human HCN2 pacemaker channel. <i>Pflugers Arch.</i> Vol. 439, No. 5, March 2000, pages 618-626;			
	40	Santoro B, et al.: Molecular and functional heterogeneity of hyperpolarization-activated pacemaker channels in the mouse CNS. <i>Jneurosci.</i> Vol. 20, No. 14, July 15, 2000, pages 5264-5275;			
	41	Valiunas, V.: Biophysical properties of connexin-45 gap junction hemichannels studied in vertebrate cells. <i>J. Gen. Physiol.</i> Vol.119, No.2, 2002, pages 147-164;			
	42	Melman YF, et al.: Structural determinants of KvLQT1 control by the KCNE family of proteins. <i>J Biol Chem.</i> Vol. 276, No. 9, March 2, 2001, pages 6439-6444;			
	43	Tinel N, et al.: KCNE2 confers background current characteristics to the cardiac KCNQ1 potassium channel. <i>EMBO J.</i> Vol. 19, No. 23, December 1, 2000, pages 6326-6330;			
	44	Martens JR, et al.: Differential targeting of Shaker-like potassium channels to lipid rafts. <i>BiolChem.</i> Vol. 275, No. 11, March 17, 2000, pages 7443-7446;			
	45	Chauhan VS, et al.: Abnormal cardiac Na(+) channel properties and QT heart rate adaptation in neonatal ankyrin(B) knockout mice. <i>Circ. Res.</i> Vol. 86, No. 4, March 3, 2000, pages 441-447;			
	46	Cao, F. et al.: A quantitative analysis of connexin-specific permeability differences of gap junctions expressed in HeLa transfectants and Xenopus oocytes. <i>J. Cell Sci.</i> Vol. 111 (pt.1), 1998, pages 31-43;			
EXAMINER			DATE CONSIDERED		
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	47	Gerhardstein BL, et al.: Proteolytic processing of the C terminus of the alpha (1C) subunit of L-type calcium channels and role of a proline-rich domain in membrane tethering of proteolytic fragments. <i>J Biol. Chem.</i> Vol. 275, No. 12, March 24, 2000, pages 8556-8563;			
	48	Barbuti A, et al.: Action of internal pronase on the f-channel kinetics in the rabbit SA node. <i>J. Physiol.</i> Vol. 520, No. 3, November 1, 1999, pages 737-744;			
	49	Wahler GM: Developmental increases in the inwardly rectifying potassium current of rat ventricular myocytes. <i>Am. J. Physiol.</i> Vol. 262, No. 5 Pt. 1, May 1992, pages C1266;			
	50	Sanguinetti MC, et al.: Coassembly of KvLGQT1 and minK (I_{SK}) proteins to form cardiac I_{SK} potassium channels. <i>Nature.</i> Vol. 384, No. 6604, November 7, 1996, pages 80-83;			
	51	Dixon JE and McKinnon D: Quantitative analysis of potassium channel expression in atrial and ventricular muscle of rats. <i>Circ. Res.</i> Vol. 75, No. 2, August 1994, pages 252-260;			
	52	Abbott GW, et al.: MiRP1 forms I_{kr} potassium channels with HERG and is associated with cardiac arrhythmia. <i>Cell.</i> Vol. 97, No. 2, April 16, 1999, pages 175-187;			
	53	Altomare C, et al.: Integrated allosteric model of voltage gating of HCN channels. <i>J.Gen. Physiol.</i> Vol. 117, No.6, 2001, pages 519-532;			
	54	Valiunas, V. et al.: Cardiac gap junction channels show quantitative differences in selectivity. <i>Circ. Res.</i> Vol.91, No.2, 2002, pages 104-111;			
	55	Accili EA, et al.: Properties and modulation of I_f in newborn versus adult cardiac SA node. <i>Am. J. Physiol.</i> Vol. 272, 1991, pages H1549-H1552;			
	56	Accili EA, et al.: Differential control of the hyperpolarization-activated current (I_f) by intracellular cAMP and phosphatase inhibition. <i>J. Physiol.</i> Vol. 491, 1996, pages 643-651;			
EXAMINER			DATE CONSIDERED		
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					

Form PTO-1449		U.S. Department of Commerce Patent and Trademark Office		Atty. Docket No. 68262-A/JPW/PJP	Serial No. 10/757,827
INFORMATION DISCLOSURE STATEMENT (Use several sheets if necessary)				Applicant Peter Brink et al.	
				Filing Date January 15, 2004	Group
OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)					
	57	Wainger BJ, et al.: Molecular mechanism of cAMP modulation of HCN pacemaker channels. <i>Nature</i> . Vol. 411, No. 6839, 2001, pages 805-810;			
	58	Ellingsen O, et al.: Adult rat ventricular myocytes cultured in defined medium: phenotype and electromechanical function. <i>Am. J. Physiol.</i> Vol. 265, No. 2 Pt. 2, August 1993, pages H747-H754.			
EXAMINER		DATE CONSIDERED			
*EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this from with next communication to applicant.					